## CLASS 11 | PHYSICS



## **CHAPTER-8** | Mechanical Properties of Solids

QUIZ PART-01

	TARTOL
<ol> <li>The property of a body to regain its original shape and size after removal of deforming force is called:         <ul> <li>A. Plasticity</li> <li>B. Elasticity</li> <li>C. Elastic fatigue</li> <li>D. Restoring force</li> </ul> </li> </ol>	<ul> <li>6. Which type of stress is produced when the length of a body increases due to applied force?</li> <li>A. Shearing stress</li> <li>B. Compressive stress</li> <li>C. Tensile stress</li> </ul>
<ul> <li>Explanation: Elasticity is defined as the ability of a body to return to its original state when the deforming force is removed.</li> <li>Which of the following is the nearest example of a perfectly elastic body? <ul> <li>A. Wax</li> <li>B. Clay</li> </ul> </li> </ul>	D. Volume stress  (C)  Explanation: Tensile stress is the longitudinal stress produced due to an increase in length.  7. If a body is subjected to equal forces from all directions, the stress experienced is:  A. Shearing stress C. Tensile stress  D. Postoring stress
<ul> <li>Explanation: Quartz is the closest natural example of a perfectly elastic body.</li> <li>3. The maximum deforming force up to which a body regains its original shape after removal of the force is known as: <ul> <li>A. Elastic fatigue</li> <li>B. Limit of elasticity</li> <li>C. Plastic limit</li> <li>D. Restoring limit</li> <li>(B)</li> </ul> </li> <li>Explanation: Beyond this limit, the body</li> </ul>	<ul> <li>D. Restoring stress</li> <li>Explanation: Force applied uniformly from all sides produces hydrostatic or volume stress.</li> <li>8. Which type of stress arises when a deforming force acts tangentially to a surface? <ul> <li>A. Longitudinal stress</li> <li>B. Volume stress</li> <li>C. Shearing stress</li> <li>D. Tensile stress (C)</li> </ul> </li> <li>Explanation: Tangential force per unit area produces shearing stress, leading to a change in shape.</li> <li>9. Strain is defined as: <ul> <li>A. Force per unit area</li> </ul> </li> </ul>
<ul> <li>cannot return to its original state.</li> <li>4. Temporary loss of elasticity caused by repeated application and removal of force is termed: <ul> <li>A. Creep</li> <li>B. Elastic fatigue</li> <li>C. Shear strain</li> <li>D. Compressive stress</li> <li>(B)</li> </ul> </li> <li>Explanation: Elastic fatigue is the temporary loss of elasticity when a force is varied frequently.</li> <li>5. Stress is mathematically expressed as: <ul> <li>A. ΔL/L</li> <li>B. F/A</li> </ul> </li> </ul>	<ul> <li>B. Ratio of change in dimension to original dimension</li> <li>C. Stress per unit strain</li> <li>D. Energy stored per unit volume</li> <li>(B)</li> <li>Explanation: Strain has no units and is given as change in dimension / original dimension.</li> <li>10. Which of the following correctly describes intermolecular forces?</li> <li>A. Always attractive</li> <li>B. Always repulsive</li> <li>C. Negligible at very large distance, attractive at intermediate distance, repulsive at very small distance</li> <li>D. Repulsive at large distances and attractive</li> </ul>
C. $\Delta V/V$ D. $\Delta \theta/\theta$ (B) Explanation: Stress is defined as the internal restoring force per unit area of crosssection.	at small distances (C)  Explanation: The behavior of intermolecular forces follows this distance-dependent pattern.